



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

12 11. 2004

Bescheinigung

Certificate

Attestation 17 DEC 2004

WIPO PCT

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03025959.2

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk



Anmeldung Nr:  
Application no.: 03025959.2  
Demande no:

Anmeldetag:  
Date of filing: 13.11.03  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Ferring B.V.  
Polaris Avenue 144  
2132 JX Hoofddorp  
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se referer à la description.)

Blister pack and solid dosage form therefor

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

A61K/

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI

BLISTER PACK AND SOLID DOSAGE FORM THEREFOR

## Field of the Invention

The present invention relates to a blister pack for pharmaceutical use comprising blisters containing a solid dosage form of desmopressin, or a pharmaceutically acceptable salt thereof, and to said solid dosage form.

## Background

Desmopressin, also known as dDAVP, is a nonapeptide and the therapeutically active ingredient (as its acetate salt) in the pharmaceutical product Minirin®, which is marketed *inter alia* as a nasal spray and a tablet formulation. Desmopressin is primarily used in the treatment of primary nocturnal enuresis, i.e. bedwetting, in children, but it is approved also for the treatment of nocturia and diabetes insipidus. The first market introduction of the tablet formulation was in Sweden in 1987. The composition of the marketed tablet form of desmopressin has remained the same to the present date.

The tablet form of desmopressin was first disclosed as set forth in the patent US 5,047,398. The subsequently issued marketing authorisations relate to a tablet where *i.a.* the mannitol, talc and cellulose components exemplified in US 5,047,398 are replaced with potato starch. In addition to desmopressin acetate and potato starch, the present tablet components are lactose, polyvinylpyrrolidone (PVP) and magnesium stearate that together form a homogeneous tablet compressed from a granulate. As a mixture of water and ethanol is used as granulation liquid in the granulate preparation, the resulting tablet also contains minor residues of those two solvents, typically 5-6% of water and 0.1% of ethanol (percentage by weight). Complete removal of residual solvents is neither required nor practical, as conditions for complete drying of solid dosage forms tend to be



Disclosure of the Invention

The present invention relates to a blister pack for pharmaceutical use comprising blisters containing a solid dosage form of desmopressin, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable adjuvant, diluent and/or carrier, wherein said solid dosage form is adapted to prevent moisture related degradation of said desmopressin.

The present solid dosage form may optionally comprise at least one further additive typically selected from a disintegrating agent, binder, lubricant, flavoring agent, preservative, colorant and any suitable mixture thereof. Examples of additives that may be considered in practising the present invention are found in "Handbook of Pharmaceutical Excipients"; Ed. A.H. Kibbe, 3<sup>rd</sup> Ed., American Pharmaceutical Association, USA and Pharmaceutical Press UK, 2000.

Without being bound by a particular theory, the inventors hypothesise that the presence of residual moisture in solid dosage forms of desmopressin in combination with the increased potential influx of moisture in blister packs (compared e.g. to sealed bottles) caused the aforementioned accelerated degradation of desmopressin upon storage. The presence of moisture in solid dosage forms appears to promote dimer formation, i.e. deactivation, of desmopressin, albeit also other variants of deactivated desmopressin are formed during storage.

More specifically it has been found that a purposive selection and control of the pH level in a solid dosage form of desmopressin is particularly efficient in counteracting degradation upon storage in blister packs.

A preferred embodiment of the present invention relates to said blister pack, wherein said solid dosage form contains an agent that provides a pH in the range of from 3.0 to 6.2 as measured when said solid dosage form

4

is contacted with water. In another embodiment said pH is in the range of from 3.0 to 6.0. As used herein the expression "contacted with water" refers to preparing a slurry of 1 g of a solid dosage form in 2 ml H<sub>2</sub>O at 25°C, where the slurry is subjected to a conventional pH measurement. A pH meter of type pH3359-9 provided by Radiometer Analytical S.A. (France) was utilised for the measurements. A slurry of 1 g of the previously known Minirin® tablet in 2 ml H<sub>2</sub>O provides a pH of about 6.6 at 25°C.

It is preferred that said pH is in the range of from 3.5 to 5.5. It is even more preferred that said pH is in the range of from 4.0 to 5.0, preferably from 4.5 to 4.8.

Said agent providing said pH is preferably an acid, preferably an acid selected from a group consisting of citric acid, hydrochloric acid and malic acid. Other examples of suitable acids are stearic acid, acetic acid, phosphoric acid, adipic acid, tartaric acid, glutamic acid and aspartic acid. The possibility of using one substance only as the pH controlling agent makes the present invention particularly convenient to practise.

In the present blister pack said blisters, and also lid foil as suitable, are preferably composed of a material selected from PVC, PVC/PVDC blends, PE (polyethylene), PP (polypropylene), polystyrene, polyester (e.g. a polyester terephthalate), paper, polyamide, PET (polyethylene terephthalate), COC (cyclic olefin copolymer) and aluminium foil or any blend thereof. As used herein the expression "blend" also encompasses a layered composite. PVC is the preferred material.

A typical aluminium blister is made of a blend which is usually a layered composite of oriented polyamide (OPA), aluminium and polypropylene, or PVC, as the bottom web, whereas the lid foil consists of aluminium. The lid foil is typically provided with a heat sealing lacquer for sealing e.g. with the polypropylene. The typical COC blister is made of PP/COC/PP as bottom web, where the

aforementioned type of aluminium foil lid may be used. The PVC blister is typically sealed with an aluminium foil lid provided with a conventional heat sealing lacquer adapted for sealing with the PVC. The blister pack  
5 manufacturing technology per se utilised in practising the present invention is conventional for a person skilled in the art. Examples of commercial providers are the Alcan Packaging Group (Singen, DE) for aluminium and COC, Riblex Film A/S (Denmark) for PVC and Perlen Con-  
10 verting AG (Switzerland) for PVDC.

In the most preferred embodiment said solid dosage form does not comprise an enteric coating. By entirely avoiding solid dosage forms with an enteric coating the preparation of the solid dosage form is simplified sub-  
15 stantially, which is a considerable practical advantage of this particular embodiment.

Said solid dosage form is preferably selected from a group consisting of tablets, granulate powder, lozenge, cachet, dry powder, wafer sheet and capsule. A tablet is  
20 most preferred.

A second aspect of the present invention relates to a solid dosage form of desmopressin, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable adjuvant, diluent and/or  
25 carrier, wherein said solid dosage form comprises an agent that provides a pH in the range of from 4.5 to 5.5 as measured when said solid dosage form is contacted with water; with the proviso that said solid dosage form does not comprise fish gelatin or an enteric coating.

30 In this second aspect it is preferred that said pH is in the range of from 4.5 to 5.0, preferably from 4.5 to 4.8. Said agent is typically an acid, preferably an acid selected from a group consisting of citric acid, hydrochloric acid and malic acid. Other examples of acids  
35 are those previously listed. The solid dosage form as such is preferably selected from a group as previously listed. Tablet is the most preferred alternative.

6

Accordingly the present invention also relates to a blister pack for pharmaceutical use comprising blisters containing a solid dosage form as defined in the second aspect of the present invention. Said blisters, and lid foil as suitable, are preferably composed of a material selected among those already listed above.

The present pharmaceutical composition in a solid dosage form is typically a perorally available tablet. A tablet may be manufactured by compression of a suitable granulate by procedures well established in the art. Some examples of suitable tablet compressing equipment are rotary presses provided by Elizabeth-Hata International, USA, and Courtoy NV, BE. For a comprehensive overview of pharmaceutical tablet manufacturing, see "Tableting" (by N.A. Armstrong) in said "Pharmaceutics - The science of dosage form design".

The following examples illustrate the present invention in more detail. They shall not be construed as a limitation of how the invention may be practised.

## Experimental

Experimental  
Example 1: Preparation of acid containing ("Low pH";  
02K24-01) solid dosage form of dDAVP

Lactose (900 g, Pharmatose 150M; provided by DMV, NL) and starch (550 g, AmylSolVät; provided by Lyckeby Stärkelse AB, SE) are mixed and sieved through a 1 mm sieve. A granulation liquid consisting of malic acid (1.88 g), water (75 ml) and PVP (13.8 g, Kollidon® 25; provided by BASF GmbH, DE) is prepared, to which dDAVP (0.75 g; provided by PolyPeptide Laboratories AB, SE) and ethanol (225 g) are added. The granulation liquid is then added to the lactose/starch mixture. After sieving (1.4 mm), drying for 20-25 hours at 40°C and further sieving (1.4 mm), the obtained granulate is admixed with magnesium stearate (11.3 g, 1.0 mm sieved; provided by Peter Greven NV, NL) and subsequently compressed to 7500 tablets using a single punch tablet compression machine



7

(Fette Exakta 1). A typical prepared tablet containing 0.1 mg of dDAVP is white, convex and oval (6.8 x 9.6 mm) with a thickness of 3-4 mm and a target weight of 192 mg. It has a smooth surface without scratches or chipped edges, and shows no tendencies to lamination (so-called capping). The residual water content is 6.1% by weight. The pH of a slurry in water of the prepared tablet is 4.6 at 25°. The pH of the dried granulate, i.e. the tablet precursor material which may also be used as a solid dosage form per se, is 4.3.

Example 2: Preparation of acid free ("Standard pH"; DK7333) solid dosage form of dDAVP

Tablets are prepared as in example 1, albeit the malic acid is omitted. The pH of a slurry in water of the prepared tablet is 6.6 at 25°. The residual water content is 6.1% by weight, i.e. intentionally identical with that of the tablet in example 1.

Example 3: Incorporation of the tablets of examples 1 and 2 in PVC blisters and stability study A

20 Blister packs are prepared using conventional packaging technology provided by Inpac AB, Lund, SE. PVC blisters (RN23-A, batch #26145-1; provided by Riblex Film A/S) and an aluminium lid foil (K7606002, batch #771297; provided by said Alcan Packaging Group were utilised  
25 together with a heat lacquer (Termolack LA723) for sealing.

The blister packages containing the tablets of examples 1 and 2 were stored at 40°C at a relative humidity (RH) of 75% in climate chambers. The content of water/moisture (% by weight) and intact dDAVP (start content 100% at 0 months) were monitored over time, and the results are summarised in Table 1.

The analytical methods used were conventional Karl Fischer and LC/UV for the water and desmopressin, respectively.



In summary, the above results show that a lower pH provides an increased storage stability for a solid dosage form of desmopressin in blister packs.

5 All references listed are to be regarded as an integral part of the present writ.

10

## CLAIMS

1. Blister pack for pharmaceutical use comprising blisters containing a solid dosage form of desmopressin, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable adjuvant, diluent or carrier, wherein said solid dosage form is adapted to prevent moisture related degradation of said desmopressin.

10            2. Blister pack according to claim 1, wherein said solid dosage form contains an agent that provides a pH in the range of from 3.0 to 6.2 as measured when said solid dosage form is contacted with water.

3. Blister pack according to claim 2, wherein said  
15 pH is in the range of from 3.5 to 5.5.

4. Blister pack according to claim 3, wherein said pH is in the range of from 4.0 to 5.0, preferably from 4.5 to 4.8.

5. Blister pack according to any one of claims 1-4,  
20 wherein said agent is an acid, preferably an acid se-  
lected from a group consisting of citric acid, hydrochlo-  
ric acid and malic acid.

6. Blister pack according to any one of claims 1-5, wherein said blisters are composed of a material selected from PVC, PVC/PVDC blends, PE, PP, polystyrene, polyester, paper, polyamide, PET, COC, aluminium foil and blends thereof.

7. Blister pack according to any one of claims 1-6, wherein said solid dosage form does not comprise an enteric coating.

8. Blister pack according to any one of claims 1-7, wherein said solid dosage form is selected from a group consisting of tablets, granulate powder, lozenge, cachet, dry powder, capsule and wafer sheet.

35 9. Solid dosage form of desmopressin, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable adjuvant, diluent or carrier,

11

wherein said solid dosage form comprises an agent that provides a pH in the range of from 4.5 to 5.5 as measured when said solid dosage form is contacted with water; with the proviso that said solid dosage form does not comprise fish gelatin or an enteric coating.

10. Solid dosage form according to claim 9, wherein said pH is in the range of from 4.5 to 5.0, preferably from 4.5 to 4.8.

11. Solid dosage form according to any one of claims 9-10, wherein said agent is an acid, preferably an acid selected from a group consisting of citric acid, hydrochloric acid and malic acid.

12. Solid dosage form according to any one of claims 9-11, which is selected from a group consisting of tablets, granulate powder, lozenge, cachet, dry powder, capsule and wafer sheet.

13. Blister pack for pharmaceutical use comprising blisters containing a solid dosage form as defined in any one of claims 9-12.

20 14. Blister pack according to claim 13, wherein said blisters are composed of a material selected from PVC, PVC/PVDC blends, PE, PP, polystyrene, polyester, paper, polyamide, PET, COC, aluminium foil and blends thereof.

